

Spine Pre Surgery orders

R3601 Rev. 5/2020 Page 1 of 1

To be performed within 30 days, unless otherwise noted.

Fax to (513) 585-0169

Surgeon name:

Phone: Fax:

Patient Name :

Date of Birth

Surgery confirmation #



Procedure Orders: _____

WEIGHT (kg): _____ **ALLERGIES:** _____

General/MAC/Regional Anesthesia **Pre Admission Testing/Same Day Surgery RN to check if below criteria is met**
ECG required - within 6 months of surgery **if:**
 Diagnosis of: CAD, arrhythmia, CHF, CRF, arterial vascular disease, pulmonary disease (except asthma), DM
PT/INR day of surgery **required** – if no documented INR of 1.1 or less within 48 hours of surgery if on Warfarin in the last 30 days
POCT Glucose day of surgery **required** – if diabetic, if blood glucose is less than 71 mg/dL (or less than 100mg/dL and symptomatic) or if greater than or equal to 180 mg/dL, initiate Preop Diabetes/Glycemic Control Order Set and Notify Anesthesia.
Potassium day of surgery **required** – **if:** 1) On dialysis or 2) Diagnosis of renal failure (not renal insufficiency or working transplant)
 Urine pregnancy (Beta HCG if unable to void) on day of surgery, unless patient has negative serum pregnancy test within 7 days of surgery – **required** if female with no history of hysterectomy **and:**
 1) 11-55 years 2) Less than 11 years and has begun menses or 3) Greater than 55 years and less than one year post-menopausal
IV: Insert Peripheral IV day of surgery (and saline lock if needed per anesthesia)
 Normal Saline @ 125 ml/hr (1000 ml bag) unless diagnosed with CRF then @ 50 ml/hr
 Other IV _____
 Local Anesthesia

Pre-operative consultation to evaluate for risk factors prior to surgery
 per PCP, may use hospitalists if not available per hospitalist per surgeon: date _____

Request for anesthesia to provide postoperative advanced pain management

Nursing Communication:
 2% Chlorhexidine Gluconate wash cloths to be used to operative site upon arrival
 Pre-Op: TED Hose Right Left Bilateral Height Knee Thigh
 Incentive Spirometer Pulmonary Function Test scheduled/results on chart
Pre Admission Testing/Same Day Surgery RN to check if below criteria is met
 Staph PCR/Culture Protocol for Staph Aureus
 If Nasal PCR/culture not performed prior to OR, send nasal PCR/culture to rule out Staph Aureus as STAT, place order for Mupirocin (Bactroban) 2%, preop once, and apply to both nostrils.
 If nasal PCR/culture not final prior to OR, order Mupirocin (Bactroban) 2%, preop once, and apply to both nostrils.
 If nasal PCR/culture obtained preoperatively, and results were positive for Staph Aureus or MRSA, record start date for Mupirocin (Bactroban) in the PTA meds. If Mupirocin (Bactroban) has not been started, order Mupirocin (Bactroban) 2%, preop once, and apply to both nostrils.
 If nasal PCR/culture positive for MRSA, discontinue current preop antimicrobial selection, and order Vancomycin 15mg/kg (max 2000mg) x1.

Labs: CBC Renal (BMP) Type & Screen PT/INR PTT CBC w/ differential Fibrinogen
 Nasal PCR or nasal culture (if on nasal steroid) for *Staph aureus* Albumin Pre-Albumin HTLV
 Carboxyhemoglobin level Urine culture Urine Cotinine levels (COT) Urinalysis _____
 Nursing Communication: HgbA1c for DIABETIC patients if not within last 30 days COVID19
 Enhanced Recovery Labs: Hemoglobin & Hematocrit, Albumin, and Prealbumin

Diagnostic Tests:
X-ray: Chest PA & Lateral (within 6 months) **Reason:** _____
EKG: 12 Lead EKG **Reason:** _____

VTE Mechanical Prophylaxis: (MUST CHOOSE ONE)
 Place SCD prior to induction of anesthesia Knee Thigh Foot Right Left Bilateral
 NO SCD needed-must give reason Already anticoagulated Ambulating Patient Refused Fall risk Not indicated-low clinical risk

VTE Pharmacologic Prophylaxis:
 Heparin 5,000 units, subcutaneous, preop once
 No pharmacologic VTE-must give reason Already anticoagulated Bleeding risk Active bleeding Patient Refused
 Thrombocytopenia Not-indicated-low clinical risk

No preop antibiotics needed

Pre-Operative Antibiotics: *Required -*Spine
 Cefazolin 2 g IVPB x1 if patient greater than or equal to 120 kg Cefazolin 3 g IVPB x1 **Alternate if allergy give** Clindamycin 900 mg IVPB x1
 History of MRSA infection Vancomycin 15 mg/kg IVPB x1 Maximum dose of 2000 mg
 Enhanced Recovery Medications: gabapentin 600 mg oral give 1.5 hours prior to surgery x1 and Oxycodone ER 10 mg oral pre-op x1

Physician Signature _____ Date: _____ Time: _____