

**Hybrid OR Cath Lab/EP Pre Procedure Orders**

R3760 Rev. 3/2022 Page 1 of 1

To be performed within 30 days, unless otherwise noted.

Fax to (513) 585-0169

Surgeon name:

Phone:

Fax:

Patient Name :

Date of Birth

Surgery confirmation #

Procedure Orders: \_\_\_\_\_

WEIGHT (kg): \_\_\_\_\_ ALLERGIES: \_\_\_\_\_

**General/MAC/Regional Anesthesia**      **Pre Admission Testing/Same Day Surgery RN to check if below criteria is met**  
**ECG required** - within 6 months of surgery **if:**  
 Diagnosis of: CAD, arrhythmia, CHF, CRF, arterial vascular disease, pulmonary disease (except asthma), or DM  
**PT/INR** day of surgery **required** – if no documented INR of 1.1 or less within 48 hours of surgery if on Warfarin in the last 30 days  
**POCT Glucose** day of surgery **required** – if diabetic, if blood glucose is less than 71 mg/dL (or less than 100mg/dL and symptomatic) or if greater than or equal to 180 mg/dL, initiate Preop Diabetes/Glycemic Control Order Set and Notify Anesthesia  
**Potassium** day of surgery **required** – **if:** 1) On dialysis or 2) Diagnosis of renal failure (not renal insufficiency or working transplant)  
**Urine pregnancy** (Beta HCG if unable to void) on day of surgery, unless patient has negative serum pregnancy test within 7 days of surgery – **required** if female with no history of hysterectomy **and:**  
 1) 11-55 years 2) Less than 11 years and has begun menses or 3) Greater than 55 years and less than one year post-menopausal  
**IV: Insert Peripheral IV day of surgery (and saline lock if needed per anesthesia)**  
 Normal Saline @ 125 ml/hr (1000 ml bag) unless diagnosed with CRF then @ 50 ml/hr  
 Other IV \_\_\_\_\_  
 **Local Anesthesia**

**Pre-operative consultation to evaluate for risk factors prior to surgery**  
 per PCP, may use hospitalists if not available     per hospitalist     per surgeon: date \_\_\_\_\_

Nursing:  Void on call to OR

**Labs:**  CBC with diff     Type & Screen     PT/INR     PTT     HgbA1C  
 Renal (BMP)     Albumin     Plasma free Hgb     Haptoglobin     CK-MB  
 Urinalysis     LDH     LIPV     Amylase     Lipase  
 COVID19     Other: \_\_\_\_\_     Other: \_\_\_\_\_

**Diagnostic Studies:**  Chest X-ray PA & Lateral     Other: \_\_\_\_\_  
**Reason:** \_\_\_\_\_  
 **ECG Reason:** \_\_\_\_\_

**VTE Mechanical Prophylaxis:**  
 Place SCD prior to induction of anesthesia     Knee     Thigh     Right     Left     Bilateral  
 NO SCD needed-must give reason     Already anticoagulated     Ambulating     Refused     Comfort measures only     Fall risk  
 Not indicated-low clinical risk

**VTE Pharmacologic Prophylaxis:**  
 Heparin 5,000 units, subcutaneous, preop once  
 No pharmacologic VTE-must give reason     Already anticoagulated     Bleeding risk     Active bleeding     Refused     Comfort measures only  
 Thrombocytopenia     Not-indicated-low clinical risk

No preop antibiotics needed

**Pre-Operative Antibiotics:**  
**\*Required: \*Pacemaker or defibrillator implant**  
 Cefazolin 2g IVPB if patient greater than or equal to 120kg Cefazolin 3g IVPB x1 **Alternate if allergy give** Vancomycin 15mg/kg (Max 2000mg) x1  
 Other: \_\_\_\_\_

Physician Signature \_\_\_\_\_ Date: \_\_\_\_\_ Time: \_\_\_\_\_

