Plastics Pre Surgery orders

R3588 Rev. 3/2022 page 1 of 1 To be performed within 30 days, unless otherwise noted.

Fax to (513) 585-0169

Surgeon name: Phone:

Fax:

Patient Name :

Date of Birth

Surgery confirmation #

Proceedings Andersi
Procedure Orders:
WEIGHT (kg): ALLERGIES:
□ General/MAC/Regional Anesthesia Pre Admission Testing/Same Day Surgery RN to check if below criteria is met ECG required - within 6 months of surgery if: Diagnosis of: CAD, arrhythmia, CHF, CRF, arterial vascular disease, pulmonary disease (except asthma), or DM PT/INR day of surgery required – if no documented INR of 1.1 or less within 48 hours of surgery if on Warfarin in the last 30 days POCT Glucose day of surgery required – if diabetic, if blood glucose is less than 71 mg/dL (or less than 100mg/dL and symptomatic) or if greater than or equal to 180 mg/dL, initiate Preop Diabetes/Glycemic Control Order Set and Notify Anesthesia Potassium day of surgery required – if: 1) On dialysis or 2) Diagnosis of renal failure (not renal insufficiency or working transplant) Urine pregnancy (Beta HCG if unable to void) on day of surgery, unless patient has negative serum pregnancy test within 7 days of surgery – required if female with no history of hysterectomy and: 1) 11-55 years 2) Less than 11 years and has begun menses or 3) Greater than 55 years and less than one year post-menopausal IV: Insert Peripheral IV day of surgery (and saline lock if needed per anesthesia) Normal Saline @ 125 ml/hr (1000 ml bag) unless diagnosed with CRF then @ 50 ml/hr □ Local Anesthesia Pre-operative consultation to evaluate for risk factors prior to surgery □ per NCP, may use hospitalists if not available
per PCP, may use nospitalists if not available per nospitalist per surgeon: date
Nursing: □ Do not give any pre-op Sedatives. The patient needs to stand up in OR room prior to surgery. □ Do not give any pre-op Sedatives until seen by MD in SDS □ Notify MD if SBP > 140 and/or DBP is > 90
Labs: CBC PT/INR PTT COVID19 Type & Screen Basic Metabolic Panel (EP1) Other:
Diagnostic Studies: ECG Reason:
VTE Mechanical Prophylaxis: Place SCD prior to induction of anesthesia Knee Thigh Right Left Bilateral NO SCD needed-must give reason Already anticoagulated Ambulating Refused Comfort measures only Fall risk Not indicated-low clinical risk
VTE Pharmacologic Prophylaxis: Heparin 5,000 units, subcutaneous, preop once No pharmacologic VTE-must give reason Already anticoagulated Bleeding risk Active bleeding Refused Comfort measures only Thrombocytopenia Not-indicated-low clinical risk
□No preop antibiotics needed
Pre-Operative Antibiotics: Cefazolin 2g IVPB x1 if patient greater than or equal to 120kg Cefazolin 3g IVPB x1 Alternate if allergy give Clindamycin 900 mg IVPB x1

Physician Signature _____

Date: _____ Time: _

