

ANTERIOR CRUCIATE LIGAMENT (ACL) AND POSTEROLATERAL CORNER (PLC) RECONSTRUCTION Physical Therapy Protocol

Patient Name:	Date of Surgery:
Evaluate and Treat – no open chain or isokinet	ic exercises
Provide patient with home exercise program	
Frequency:x/week xweeks	
Phase I (0-6 weeks):	
Weight Bearing: Heel touch WB in brace.*	
Brace: 0-2 week: Locked in full extension for ambulation 2.6 week by the land for a week the firm 0.00 many and the second	
2-6 weeks: Unlocked for ambulation 0-90, remov	e for sleeping.**
• ROM: 0-2 week: 0- 45.	
2-6 weeks: Advance slowly 0-90.	
• Exercises: Quad sets, patellar mobs, gastroc/soled SLR w/ brace in full extension until qua Side-lying hip/core. Hamstrings avoidance until 6 weeks po	d strength prevents extension lag.
Phase II (6-12 weeks):	
• Weight Bearing: Advance 25% weekly until full b	y 8 weeks.
• Brace: Discontinue at 6 wks if no extension lag.	
• ROM: Full.	
 Exercises: Begin toe raises, closed chain quads, be step-ups, front and side planks; advance 	· · · · · · · · · · · · · · · · · · ·
Phase III (12-16 weeks):	
• Weight Bearing: Full.	
• Brace: None.	
• ROM: Full.	
• Exercises: Advance closed chain strengthening.	

Progress proprioception activities.

ahead at 12 weeks.

Begin stairmaster, elliptical and running straight



Phase IV (16-24 weeks):	
 Weight Bearing: Full. 	
• Brace: None.	
• ROM: Full.	
 Exercises: 16 weeks: Begin jumping . 20 weeks: Advance to sprinting, backward running program and sport-specific drills. 	, cutting/pivoting/changing direction, initiate plyometric
Phase V (>6 months):	
• Weight Bearing: Full.	
• Brace: None.	
• ROM: Full and pain-free.	
• Exercises: Gradual return to sports participation af Maintenance program based on FSA.	ter completion of FSA.***
*Modified with concomitantly performed meniscus repair/transplanta	ation or articular cartilage procedure
**Brace may be removed for sleeping after first post-operative visit (da	ay 7-10)
***Completion of FSA (Functional Sports Assessment) not mandatory, for competitive athletes returning to play after rehab	, but recommended at 22-24 wks post-op
By signing this referral, I certify that I have examined this p This patient would would not benefit from	
Physician Name:	Date:

